

Table S5. Estimated v and correction factor for each substrate.

Substrate	estimated v (pmol/min/mg)				
	[S] ; 0.1 μ M	[S] ; 1 μ M	[S] ; 2 μ M	[S] ; 25 μ M	[S] ; 250 μ M
Taurocholate	0.301 (0.103)	2.91 (1)	5.60 (1.92)	38.1 (13.1)	69.9 (24.0)
Estrone-3-sulfate	0.650 (0.107)	6.06 (1)	11.3 (1.86)	55.5 (9.16)	80.1 (13.2)
Cholecystokinin Octapeptide	2.60 (0.123)	21.2 (1)	35.1 (1.66)	88.5 (4.17)	100 (4.72)
E2-17G	0.152 (0.104)	1.46 (1)	2.81 (1.92)	18.5 (12.7)	33.0 (22.6)
MPP ⁺	2.24 (0.109)	20.6 (1)	37.6 (1.83)	159 (7.73)	212 (10.3)
Digoxin	0.180 (0.136)	1.32 (1)	2.05 (1.55)	4.10 (3.11)	4.45 (3.37)

E2-17G, Estradiol 17 β -(D)-glucuronide.

The numbers shown in parentheses represents correction factors for normalization to estimate corrected velocities (V_{max}) at 1 μ M substrate concentration.