

Original Article

Residual metals in carbon nanotubes suppress the proliferation of neural stem cells

Yukari Shigemoto-Mogami¹, Koki Fujimori^{1,2}, Yoshiaki Ikarashi³, Akihiko Hirose⁴,
Yuko Sekino¹ and Kaoru Sato¹

¹Laboratory of Neuropharmacology, Division of Pharmacology, National Institute of Health Sciences,
1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

²Division of Basic Biological Science, Faculty of Pharmacy, Keio University,
1-5-30 Shiba-koen, Minato-ku, Tokyo 105-8512, Japan

³Division of Environmental Chemistry, National Institute of Health Sciences,
1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

⁴Division of Risk Assessment, National Institute of Health Sciences,
1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

(Received October 17, 2014; Accepted October 20, 2014)

ABSTRACT — Carbon nanotubes (CNTs) are used in many fields; however, little is known about the effects of CNTs on the central nervous system (CNS). In this study, we found that extracts of sonicated CNTs suppressed the proliferation of neural stem cells (NSCs). Single-walled CNTs (SWCNTs) and multiple-walled CNTs (MWCNTs) were suspended in PBS (1 mg/mL) and sonicated for 5 hr using a water bath sonicator. Supernatants from both types of CNTs suppressed NSC proliferation. The effects weakened in a dilution-ratio-dependent manner and strengthened in a sonication time-dependent manner. Metal concentrations extracted from SCNTs and MCNTs after 5-hr of sonication were determined using inductively coupled plasma mass spectrometry. Mn, Rb, Cs, Tl, and Fe were detected in the SWCNT supernatant, and Mn, Cs, W, and Tl were detected in the MWCNT supernatant. The concentration of Mn, Rb, and Fe eluted from the SWCNTs and Rb eluted from MWCNTs following sonication were sufficient to suppress NSC proliferation alone. N-acetyl cysteine (NAC) and ascorbic acid (AA) reversed the effects of Mn and Fe and restored NSC proliferation. The effects of Rb and Tl were not affected by the antioxidants. Both antioxidants largely restored the suppression of NSC proliferation induced by the SWCNT and MWCNT supernatants. These results suggest that metals extracted from CNTs via a strong vibration energy can suppress NSC proliferation through ROS production by the extracted metals.

Key words: Carbon nanotube, Neural stem cell, Metals, Proliferation

INTRODUCTION

CNTs are fiber-shaped nanomaterials that consist of graphite hexagonal-mesh planes (graphene sheet) in a single-layer (single-walled carbon nanotubes (SWCNTs)) or in multiple layers with nest accumulation (multi-walled carbon nanotubes (MWCNTs)). The structure of SWCNTs is a honeycomb carbon lattice rolled into a cylinder, and the basic morphology consists of a sheet of tangled SWCNT (with a diameter of approximately 2 nm) bundles with diameters tens of nanometers in length. The structure of MWCNTs consists of honeycomb carbon lattices rolled into a multi-layer tubular shape, and the basic morpho-

gy is composed of particles of tangled MWCNTs with a diameter of approximately 30 nm. CNTs are used in many fields, including energy, healthcare, environment, materials, and electronics. However, adverse effects of CNTs on human health are poorly understood. Exposure to asbestos is known to cause asbestosis, bronchogenic carcinoma, mesothelioma, pleural fibrosis and pleural plaques, indicating that both the lungs and the pleura are targets of asbestos (Donaldson *et al.*, 2013). CNTs also exist as fibers or compact particles; thus, most studies concerning the adverse effects of CNTs have focused on lung toxicity (Jaurand *et al.*, 2009; Pacurari *et al.*, 2010) based on the fiber pathogenicity paradigm developed in the 1970-80s.

Correspondence: Kaoru Sato (E-mail: kasato@nihs.go.jp)

However, recent reports showed that nano-particles can cross the blood–brain barrier (BBB) and enter the brain (Sharma and Sharma, 2007). Furthermore, it has been suggested that the olfactory nerve pathway is a portal of entry into the CNS (Henriksson and Tjalve, 2000; Persson *et al.*, 2003; Mistry *et al.*, 2009; Balasubramanian *et al.*, 2013). Recent reports showed that MWCNTs are toxic to neural cells (Belyanskaya *et al.*, 2009; Xu *et al.*, 2009; Gavello *et al.*, 2012). Here, we investigated the effects of CNTs on the self-renewal of neural stem cells (NSCs). The mammalian CNS comprises various cell types, including neurons, astrocytes, and oligodendrocytes, and these cells differentiate from NSCs at specific brain developmental stages. Sufficient proliferation of NSCs before differentiation is essential to supply the neurons and glia required for brain function (Caviness *et al.*, 1995; Kriegstein and Alvarez-Buylla, 2009). In addition, NSCs are maintained in the subventricular zone and the hippocampal subgranular zone in the adult brain. Adult neurogenesis from these NSCs plays a key role in higher-order brain functions, such as cognition, learning and memory (Couillard-Despres *et al.*, 2011; Eisch and Petrik, 2012; Rolando and Taylor, 2014). Thus, the effects of CNTs on the proliferation of NSCs need to be determined for both of brain development and brain function. Here, we report that sonicated extracts of CNTs suppressed the proliferation of NSCs. We also determined that these effects were mediated through ROS produced by residual metals in the CNTs.

MATERIALS AND METHODS

Materials

CNTs (SWCNT: purity > 95%; Lot No.: SW1859; MWCNT: purity: > 98%; Lot No.: 04-12/10#1-(4)) were supplied by Nikkiso Co., Ltd. (Shizuoka, Japan). Both test materials were not coated or modified. The detailed physicochemical properties of Nikkiso CNTs have been previously reported (Ema *et al.*, 2011; Matsumoto *et al.*, 2012). Epidermal growth factor (EGF), MnCl₂, RbCl, TiCl₃, FeCl₂, FeCl₃, and NAC were purchased from Sigma (St. Louis, Mo, USA). Fibroblast growth factor 2 (FGF2) was purchased from PeproTech (Rocky Hill, NJ, USA). AA was purchased from WAKO (Osaka, Japan). The BrdU cell proliferation assay kit was purchased from Merck (Darmstadt, Germany). B27 supplement, TrypLE Select, FBS, and DMEM were purchased from Life Technologies (Grand Island, NY, USA).

Preparation of supernatants of sonicated CNT solutions

SWCNTs and MWCNTs were suspended in PBS (1 mg/mL) and sonicated for 10 min or 5 hr using a water bath-sonicator (Hitachi-Kokusai Electric Inc., Tokyo, Japan) at a frequency of 36 kHz and a watt density of 65 W/264 cm². The supernatants of sonicated CNT suspensions were diluted with culture medium 10- to 1,000-fold.

Rat neural stem cell (NSC) culture

Rat NSCs were cultured as previously described (Reynolds *et al.*, 1992; Hamanoue *et al.*, 2009) with slight modifications. Briefly, the telencephalons were dissected from embryonic day 16 (E16) rats of either sex in ice-cold DMEM/F12. The tissue was then minced and dispersed into single cells by pipetting. Cells were then cultured in DMEM/F12 containing B27 supplement (1/200), 20 ng/mL fibroblast growth factor 2 (FGF2) and 20 ng/mL epidermal growth factor (EGF) for 7 days. The primary neurospheres were incubated with TrypLE Select for 15 min and dissociated by pipetting. Single cells were seeded in 96-well plates for the proliferation assay.

Measurement of metal concentrations

CNTs were suspended in PBS (1 mg/mL) and sonicated for 5 hr using a water bath sonicator. The metal concentrations in the CNT supernatants were quantified using an inductively coupled plasma mass spectrometer (ICP-MS) (Agilent 7500ce ICP-MS, Agilent Technologies, Santa Clara, CA, USA) fitted with a collision/reaction cell in helium mode. We first detected metals at concentrations exceeding the detection limits using a semi-quantitative analysis. Next, we determined the concentration of the detected metals (i.e., Mn, Fe, Rb, Cs, W, and Ti) using a full quantitative analysis with calibration curves.

Treatment of NSCs with the supernatants of sonicated CNT suspensions, metals, and antioxidants

NSCs were treated with the supernatants of sonicated CNT suspensions, MnCl₂ (1-100 ppb), RbCl (1-100 ppb), TiCl₃ (0.1-10 ppb), FeCl₂ (100-10,000 ppb) or FeCl₃ (100-10,000 ppb) with or without 10 μM N-acetyl cysteine (NAC) or 10 μM ascorbic acid (AA) for 24 hr.

NSC proliferation assay

We quantified NSC proliferation according the instructions from the BrdU cell proliferation assay kit (Calbiochem, Hayward, CA, USA). The primary neurospheres were dissociated into single cells and seeded in 96-well plates at a density of 2 x 10⁴ cells/

well. BrdU was added to the medium during the treatment of NSCs. After incubation, the cells were fixed, and BrdU-immuno-labeling was performed. The fluorescence intensities were used as a marker of proliferation. The fluorescence was measured at an excitation wavelength of 320 nm and emission wavelength of 460 nm with a fluorescence microplate reader (Spectra Max Microplate reader, Molecular Devices, Sunnyvale, CA, USA).

Data analysis and statistics

All data are shown as the mean \pm S.E.M. The statistical analysis was performed using Student's *t*-test or an ANOVA followed by a Tukey's test. Differences were considered to be significant at $p < 0.05$.

RESULTS

SWCNTs and MWCNTs were suspended in PBS (1 mg/mL) and sonicated for 5 hr using a water bath sonicator. The supernatants of the sonicated CNT suspensions were collected and diluted with culture medium 10- to 1,000-fold. We found that a 24-hr treatment with supernatants of SWCNT and MWCNT suppressed NSC proliferation in a dilution ratio-dependent manner (Fig. 1). The suppression of proliferation was stronger with the SWCNT supernatant when compared with the MWCNT supernatant. The effects of sonication time were also assessed. The suppressive effects of both supernatants disappeared when the sonication time was changed from 5 hr to 10 min (Fig. 2). These results suggest that the suppression of NSC proliferation is due to factors released from CNTs in a sonication time-dependent manner.

CNTs are manufactured using metallic catalysts (Ding *et al.*, 2008; Yazyev and Pasquarello, 2008; Banhart, 2009; Tyagi *et al.*, 2011). Thus, we speculated that residual metals extracted from CNTs during the 5-hr sonication may be responsible for the suppression of NSC proliferation. We therefore quantified the metal contents in the CNT supernatants. The metals in the SWCNT and MWCNT supernatants were first analyzed using ICP-MS in a semi-quantitative mode. Next, the concentrations of metals were determined using calibration curves (Table 1). We found that a 5-hr sonication induced the extraction of multiple metals from the CNTs. Mn, Rb, Cs, Tl, and Fe were detected in the SWCNT supernatant, whereas Mn, Cs, W, and Tl were detected in the MWCNT supernatant. Among these metals, the concentration of Fe in SWCNT supernatant was remarkably high (from N.D. to 7,110 ppb). The concentrations of these metals in PBS were largely negligible and did not change after a




Fig. 1. Effects of the supernatants of sonicated CNT suspensions on the proliferation of rat NSCs. The supernatants of SWCNTs and MWCNTs suppressed NSC proliferation in a dilution ratio-dependent manner. *: $p < 0.05$, **: $p < 0.01$ vs. control group ($N = 6$), ANOVA followed by a Tukey's test.




Fig. 2. Sonication time-dependence of CNT supernatant effects. The effects of SWCNT and MWCNT supernatants disappeared with a sonication time of 10 min. However, a 5-hr sonication time produced a significant suppression of NSC proliferation. *: $p < 0.05$ vs. control group ($N = 6$), ANOVA followed by a Tukey's test.

5-hr sonication.

Next, we examined the direct effects of the metals at concentration ranges detected in the supernatants. Fig. 3 shows the metals that had a suppressive effect on NSC

Table 1. Metals eluted from CNTs by sonication for 5 hr.

	Concentrations of metals (ppb) 1 ppb = 10 ⁻⁸ %					
	PBS		SWCNT		MWCNT	
sonication	-	+	-	+	-	+
Mn	nd	nd	0.33	16.04	nd	0.26
Rb	3.97	3.84	6.88	13.33	4.06	4.61
Cs	nd	nd	0.1	0.32	nd	0.59
W	nd	0.05	nd	0.08	nd	0.4
Tl	md	nd	0.05	0.17	nd	0.37
Fe	nd	nd	nd	7110	nd	nd

The metal concentrations in the supernatant of SWCNT and MWCNT were quantified using ICP-MS in a semi-quantitative mode followed by a full quantitative mode. Mn, Rb, Cs, W, Tl, and Fe were detected in the SWCNT supernatant. Mn, Rb, Cs, W, Tl, and Fe were detected in the MWCNT supernatant. The concentration of Fe in the SWCNT supernatant was remarkably high (7,110 ppb).




Fig. 3. The direct effect of metals in CNT supernatants. Mn²⁺, Rb⁺, Tl³⁺, Fe²⁺, and Fe³⁺ suppressed NSC proliferation in a concentration-dependent manner. *: p < 0.05, **: p < 0.01 vs. control group (N = 12), ANOVA followed by a Tukey's test.

proliferation (Fig. 3). Mn²⁺, Rb⁺, Tl³⁺, Fe²⁺, and Fe³⁺ suppressed the proliferation of NSCs in a concentration-dependent manner. These results indicate that Mn, Rb, and Fe were present in the SWCNT supernatant at a concentration high enough to suppress NSC proliferation. This effect was induced by the Rb in the MWCNT supernatant. Some metals are known to produce reactive oxygen species (Ding *et al.*, 2008) that can result in oxidative stress on lipids, DNA and proteins (Henriksson and Tjalve, 2000; Choi *et al.*, 2007; Alekseenko *et al.*, 2008; Kim *et al.*, 2011; Latronico *et al.*, 2013; Roth and Eichhorn, 2013; Sripetchwandee *et al.*, 2013). Thus, we examined the involvement of ROS in the suppression of NSC proliferation. N-acetyl cysteine (NAC) (10 μM) and ascorbic

acid (AA) (10 μM) are typical antioxidants that can significantly restore the suppression of the NSC proliferation caused by Mn²⁺, Fe²⁺, and Fe³⁺ (Fig. 4A). The effect of Rb and Tl were not affected by NAC or AA (data not shown). These results suggest that ROS is involved in the suppressive effects produced by Mn and Fe. We also examined whether ROS played a role in the suppression of NSC proliferation by the CNT supernatants (Fig. 4B). Both NAC and AA markedly restored the decrease in NSC proliferation caused by the SWCNT and MWCNT supernatants. We confirmed that both of these antioxidants alone did not affect NSC proliferation (data not shown). Taken together, these results suggest that the suppressive effects of the sonicated extract of CNTs were mainly caused by

Effects of residual metals in carbon nanotubes on neural stem cells




Fig. 4. Antioxidants attenuated the reduction in NSC proliferation caused by metals and CNT supernatants. The suppression of the NSC proliferation caused by Mn²⁺, Fe²⁺, Fe³⁺ (A) and the supernatants of CNTs (B) was significantly restored by NAC (10 μ M) and AA (10 μ M). *: $p < 0.05$, **: $p < 0.01$ vs. control group, #: $p < 0.05$, ##: $p < 0.01$ vs. metal or CNT-supernatant-treated groups ($N = 7$), ANOVA followed by a Tukey's test.

ROS produced by residual metals.

DISCUSSION

We found that the supernatants of sonicated CNT suspensions suppress NSC proliferation. We also determined that these effects were largely mediated by ROS production from residual metals. To demonstrate the involvement of ROS, we used the two antioxidants NAC and AA. NAC exerts its protective by increasing glutathione

levels (Yim *et al.*, 1994; Arfsten *et al.*, 2007; Li *et al.*, 2009), directly scavenging ROS, and activating ERK1/2 (Zhang *et al.*, 2011). AA is a powerful water-soluble antioxidant that acts by scavenging ROS and reactive nitrogen species (Carr and Frei, 1999; Kojo, 2004). The concentrations of NAC and AA used in this study were at a level shown to suppress the effects of ROS in previous studies (Carr and Frei, 1999; De la Fuente and Victor, 2001; Nakajima *et al.*, 2009).

Proliferative NSCs have a high endogenous ROS lev-

el (Le Belle *et al.*, 2011), and redox balance is important to regulate NSC/neural progenitor cell (NPC)-self-renewal and differentiation (Smith *et al.*, 2000; Li *et al.*, 2007; Hou *et al.*, 2012; Topchiy *et al.*, 2013). For example, mitochondrial superoxide negatively regulates NPC-self-renewal in the developmental cerebral cortex (Hou *et al.*, 2012). High levels of ROS inhibit O-2A progenitor proliferation (Smith *et al.*, 2000; Li *et al.*, 2007). In other cases, NADPH oxidase (Nox) 4-generated superoxide drives mouse NSC proliferation (Topchiy *et al.*, 2013). Ketamine-induced ROS enhanced the proliferation of NSCs derived from human embryonic stem cells (Bai *et al.*, 2013). The effect of ROS on NSC/NPC proliferation may change depending on the subcellular localization of the ROS generation and the timing of the ROS generation.

The suppression of NSC proliferation by the supernatants of both CNTs were virtually restored by the antioxidants, suggesting that the effects of CNT-supernatants were mediated through ROS stress. After a 5-hr sonication, multiple metals were detected in the SWCNT and MWCNT supernatants using ICP-MS. Mn, Rb, Cs, Tl, and Fe were detected in the SWCNT supernatant, and Mn, Cs, W, and Tl were detected in the MWCNT supernatant. Out of these SWCNT metals, the effects of Mn and Fe were reversed by antioxidants, suggesting that Mn and Fe play the main role in the suppression of NSC proliferation by CNT supernatants. In the MWCNT supernatant, the concentrations of Mn and Fe were insufficient to suppress NSC proliferation. Thus, a combination of ROS produced by multiple metals might produce synergistic suppressive effects.

Fe is essential for biological processes, but it is also known to be toxic in excess. Fe²⁺ overload into the cells and shuttling of Fe²⁺ to Fe³⁺ leads to cellular malfunctions due to ROS production (Halliwell and Gutteridge, 1992; Touati, 2000). Although Fe³⁺ has been largely considered as non-cytotoxic (Braun, 1997; Bruins *et al.*, 2000), it has its own mechanisms that can alter cell viability (Chamngpol *et al.*, 2002). Fe³⁺ shows ROS production even while bound to proteins (Alekseenko *et al.*, 2008). GSH revealed pro-oxidant effects in the presence of an exogenous Fe³⁺ (Zager and Burkhart, 1998). Furthermore, Fe²⁺ and Fe³⁺ were shown to enter brain mitochondria and cause mitochondrial depolarization and ROS production (Sripetchwadee *et al.*, 2013). Mn is also essential for biological processes, but it has been known to be a neurotoxicant in excess. Mn induces oxidative stress (Choi *et al.*, 2007; Park and Park, 2010) and the release of cytokines (Park and Park, 2010). Mn further potentiates inflammation by the release of MMP9 through ROS production

and modulation of ERK (Latronico *et al.*, 2013). Rb was also detected in the supernatants of SWCNT and MWCNT. Here, we found that Rb alone suppressed NSC proliferation in a ROS-independent manner. Rb has long been considered as nontoxic. Rb is generally used as a medical contrast medium because of its long half-life. Thus, the mechanism behind the Rb effects should be clarified quickly.

Most commercial CNTs contain ultrafine metal particles composed of Fe, Ni, Y, Co, Pb, and Cu that are used as catalysts (Ding *et al.*, 2008; Yazyev and Pasquarello, 2008; Banhart, 2009; Tyagi *et al.*, 2011). Recent studies showed that metal impurities play a major role in CNT cytotoxicity (Liu *et al.*, 2008; Kim *et al.*, 2010). The residual metals can remain in the contact solvent or embed inside the CNTs (Pumera, 2007; Fubini *et al.*, 2011; Aldieri *et al.*, 2013). In our study, the content of Fe in SWCNT was remarkable. A SWCNT is a graphene sheet protected metal core/shell of nanoparticles (Pumera, 2007). This structure may have caused the higher levels of metal impurities when compared with MWCNTs. Our data suggest that the residual metallic catalysts are released by vibration energy with a sonication frequency of 36 kHz, watt density of 65 W/264 cm² and sonication time of 5 hr. Pumera *et al.* indicated that washing with concentrated nitric acid removed up to 88% (w/w) of metal catalyst nanoparticles (Pumera, 2007). For public health and the safer applications of CNTs in nano-medicine, it is preferable to decrease the amount of the metal impurities by improving the washing process.

Conflict of interest---- The authors declare that there is no conflict of interest.

REFERENCES

- Aldieri, E., Fenoglio, I., Cesano, F., Gazzano, E., Gulino, G., Scarano, D., Attanasio, A., Mazzucco, G., Ghigo, D. and Fubini, B. (2013): The role of iron impurities in the toxic effects exerted by short multiwalled carbon nanotubes (MWCNT) in murine alveolar macrophages. *J. Toxicol. Environ. Health Part A*, **76**, 1056-1071.
- Alekseenko, A.V., Waseem, T.V. and Fedorovich, S.V. (2008): Ferritin, a protein containing iron nanoparticles, induces reactive oxygen species formation and inhibits glutamate uptake in rat brain synaptosomes. *Brain Res.*, **1241**, 193-200.
- Arfsten, D.P., Johnson, E.W., Wilfong, E.R., Jung, A.E. and Bobb, A.J. (2007): Distribution of radio-labeled N-Acetyl-L-Cysteine in Sprague-Dawley rats and its effect on glutathione metabolism following single and repeat dosing by oral gavage. *Cutan. Ocul. Toxicol.*, **26**, 113-134.
- Bai, X., Yan, Y., Canfield, S., Muravyeva, M.Y., Kikuchi, C., Zaja, I., Corbett, J.A. and Bosnjak, Z.J. (2013): Ketamine enhances human neural stem cell proliferation and induces neuronal appo-

Effects of residual metals in carbon nanotubes on neural stem cells

- tosis via reactive oxygen species-mediated mitochondrial pathway. *Anesth. Analg.*, **116**, 869-880.
- Balasubramanian, S.K., Poh, K.W., Ong, C.N., Kreyling, W.G., Ong, W.Y. and Yu, L.E. (2013): The effect of primary particle size on biodistribution of inhaled gold nano-agglomerates. *Biomaterials*, **34**, 5439-5452.
- Banhart, F. (2009): Interactions between metals and carbon nanotubes: at the interface between old and new materials. *Nanoscale*, **1**, 201-213.
- Belyanskaya, L., Weigel, S., Hirsch, C., Tobler, U., Krug, H.F. and Wick, P. (2009): Effects of carbon nanotubes on primary neurons and glial cells. *Neurotoxicology*, **30**, 702-711.
- Braun, V. (1997): Avoidance of iron toxicity through regulation of bacterial iron transport. *Biol. Chem.*, **378**, 779-786.
- Bruins, M.R., Kapil, S. and Oehme, F.W. (2000): Microbial resistance to metals in the environment. *Ecotoxicol. Environ. Saf.*, **45**, 198-207.
- Carr, A. and Frei, B. (1999): Does vitamin C act as a pro-oxidant under physiological conditions? *FASEB J.*, **13**, 1007-1024.
- Caviness, V.S.Jr., Takahashi, T. and Nowakowski, R.S. (1995): Numbers, time and neocortical neurogenesis: a general developmental and evolutionary model. *Trends Neurosci.*, **18**, 379-383.
- Chamngpol, S., Dodson, W., Cromie, M.J., Harris, Z.L. and Groisman, E.A. (2002): Fe(III)-mediated cellular toxicity. *Mol. Microbiol.*, **45**, 711-719.
- Choi, C.J., Anantharam, V., Saetveit, N.J., Houk, R.S., Kanthasamy, A. and Kanthasamy, A.G. (2007): Normal cellular prion protein protects against manganese-induced oxidative stress and apoptotic cell death. *Toxicol. Sci.*, **98**, 495-509.
- Couillard-Despres, S., Iglseider, B. and Aigner, L. (2011): Neurogenesis, cellular plasticity and cognition: the impact of stem cells in the adult and aging brain--a mini-review. *Gerontology*, **57**, 559-564.
- De la Fuente, M. and Victor, V.M. (2001): Ascorbic acid and N-acetylcysteine improve *in vitro* the function of lymphocytes from mice with endotoxin-induced oxidative stress. *Free Radic. Res.*, **35**, 73-84.
- Ding, F., Larsson, P., Larsson, J.A., Ahuja, R., Duan, H., Rosen, A. and Bolton, K. (2008): The importance of strong carbon-metal adhesion for catalytic nucleation of single-walled carbon nanotubes. *Nano Lett.*, **8**, 463-468.
- Donaldson, K., Poland, C.A., Murphy, F.A., MacFarlane, M., Chernova, T. and Schinwald, A. (2013): Pulmonary toxicity of carbon nanotubes and asbestos - similarities and differences. *Adv. Drug Deliv. Rev.*, **65**, 2078-2086.
- Eisch, A.J. and Petrik, D. (2012): Depression and hippocampal neurogenesis: a road to remission? *Science*, **338**, 72-75.
- Ema, M., Matsuda, A., Kobayashi, N., Naya, M. and Nakanishi, J. (2011): Evaluation of dermal and eye irritation and skin sensitization due to carbon nanotubes. *Regul. Toxicol. Pharmacol.*, **61**, 276-281.
- Fubini, B., Fenoglio, I., Tomatis, M. and Turci, F. (2011): Effect of chemical composition and state of the surface on the toxic response to high aspect ratio nanomaterials. *Nanomedicine (Lond)*, **6**, 899-920.
- Gavello, D., Vandael, D.H., Cesa, R., Premoselli, F., Marcantoni, A., Cesano, F., Scarano, D., Fubini, B., Carbone, E., Fenoglio, I. and Carabelli, V. (2012): Altered excitability of cultured chromaffin cells following exposure to multi-walled carbon nanotubes. *Neurotoxicology*, **6**, 47-60.
- Halliwell, B. and Gutteridge, J.M. (1992): Biologically relevant metal ion-dependent hydroxyl radical generation. An update. *FEBS Lett.*, **307**, 108-112.
- Hamanoue, M., Matsuzaki, Y., Sato, K., Okano, H.J., Shibata, S., Sato, I., Suzuki, S., Ogawara, M., Takamatsu, K. and Okano, H. (2009): Cell surface N-glycans mediated isolation of mouse neural stem cells. *J. Neurochem.*, **110**, 1575-1584.
- Henriksson, J. and Tjalve, H. (2000): Manganese taken up into the CNS via the olfactory pathway in rats affects astrocytes. *Toxicol. Sci.*, **55**, 392-398.
- Hou, Y., Ouyang, X., Wan, R., Cheng, H., Mattson, M.P. and Cheng, A. (2012): Mitochondrial superoxide production negatively regulates neural progenitor proliferation and cerebral cortical development. *Stem Cells*, **30**, 2535-2547.
- Jaurand, M.C., Renier, A. and Daubriac, J. (2009): Mesothelioma: Do asbestos and carbon nanotubes pose the same health risk? *Part. Fibre Toxicol.*, **6**, 16.
- Kim, J.E., Lim, H.T., Minai-Tehrani, A., Kwon, J.T., Shin, J.Y., Woo, C.G., Choi, M., Baek, J., Jeong, D.H., Ha, Y.C., Chae, C.H., Song, K.S., Ahn, K.H., Lee, J.H., Sung, H.J., Yu, I.J., Beck, G.R.Jr. and Cho, M.H. (2010): Toxicity and clearance of intratracheally administered multiwalled carbon nanotubes from murine lung. *J. Toxicol. Environ. Health Part A*, **73**, 1530-1543.
- Kim, K.K., Singh, R.K., Strongin, R.M., Moore, R.G., Brard, L. and Lange, T.S. (2011): Organometallic iron(III)-salophene exerts cytotoxic properties in neuroblastoma cells via MAPK activation and ROS generation. *PLoS One*, **6**, e19049.
- Kojo, S. (2004): Vitamin C: basic metabolism and its function as an index of oxidative stress. *Curr. Med. Chem.*, **11**, 1041-1064.
- Kriegstein, A. and Alvarez-Buylla, A. (2009): The glial nature of embryonic and adult neural stem cells. *Ann. Rev. Neurosci.*, **32**, 149-184.
- Latronico, T., Brana, M.T., Merra, E., Fasano, A., Di Bari, G., Casalino, E. and Liuzzi, G.M. (2013): Impact of manganese neurotoxicity on MMP-9 production and superoxide dismutase activity in rat primary astrocytes. Effect of resveratrol and therapeutic implications for the treatment of CNS diseases. *Toxicol. Sci.*, **135**, 218-228.
- Le Belle, J.E., Orozco, N.M., Paucar, A.A., Saxe, J.P., Mottahedeh, J., Pyle, A.D., Wu, H. and Kornblum, H.I. (2011): Proliferative neural stem cells have high endogenous ROS levels that regulate self-renewal and neurogenesis in a PI3K/Akt-dependant manner. *Cell Stem Cell*, **8**, 59-71.
- Li, W., Nie, S., Yu, Q. and Xie, M. (2009): (-)-Epigallocatechin-3-gallate induces apoptosis of human hepatoma cells by mitochondrial pathways related to reactive oxygen species. *J. Agric. Food Chem.*, **57**, 6685-6691.
- Li, Z., Dong, T., Proschel, C. and Noble, M. (2007): Chemically diverse toxicants converge on Fyn and c-Cbl to disrupt precursor cell function. *PLoS Biol.*, **5**, e35.
- Liu, X., Guo, L., Morris, D., Kane, A.B. and Hurt, R.H. (2008): Targeted Removal of Bioavailable Metal as a Detoxification Strategy for Carbon Nanotubes. *Carbon*, **46**, 489-500.
- Matsumoto, M., Serizawa, H., Sunaga, M., Kato, H., Takahashi, M., Hirata-Koizumi, M., Ono, A., Kamata, E. and Hirose, A. (2012): No toxicological effects on acute and repeated oral gavage doses of single-wall or multi-wall carbon nanotube in rats. *J. Toxicol. Sci.*, **37**, 463-474.
- Mistry, A., Stolnik, S. and Illum, L. (2009): Nanoparticles for direct nose-to-brain delivery of drugs. *Int. J. Pharm.*, **379**, 146-157.
- Nakajima, Y., Tsuruma, K., Shimazawa, M., Mishima, S. and Hara, H. (2009): Comparison of bee products based on assays of anti-oxidant capacities. *BMC Complement. Altern. Med.*, **9**, 4.

- Pacurari, M., Castranova, V. and Vallyathan, V. (2010): Single- and multi-wall carbon nanotubes versus asbestos: are the carbon nanotubes a new health risk to humans? *J. Toxicol. Environ. Health A.*, **73**, 378-395.
- Park, E.J. and Park, K. (2010): Induction of oxidative stress and inflammatory cytokines by manganese chloride in cultured T98G cells, human brain glioblastoma cell line. *Toxicol. In Vitro*, **24**, 472-479.
- Persson, E., Henriksson, J. and Tjalve, H. (2003): Uptake of cobalt from the nasal mucosa into the brain via olfactory pathways in rats. *Toxicol. Lett.*, **145**, 19-27.
- Pumera, M. (2007): Carbon nanotubes contain residual metal catalyst nanoparticles even after washing with nitric acid at elevated temperature because these metal nanoparticles are sheathed by several graphene sheets. *Langmuir*, **23**, 6453-6458.
- Reynolds, B.A., Tetzlaff, W. and Weiss, S. (1992): A multipotent EGF-responsive striatal embryonic progenitor cell produces neurons and astrocytes. *J. Neurosci.*, **12**, 4565-4574.
- Rolando, C. and Taylor, V. (2014): Neural stem cell of the hippocampus: development, physiology regulation, and dysfunction in disease. *Curr Top Dev Biol.*, **107**, 183-206.
- Roth, J.A. and Eichhorn, M. (2013): Down-regulation of LRRK2 in control and DAT transfected HEK cells increases manganese-induced oxidative stress and cell toxicity. *Neurotoxicology*, **37**, 100-107.
- Sharma, H.S. and Sharma, A. (2007): Nanoparticles aggravate heat stress induced cognitive deficits, blood-brain barrier disruption, edema formation and brain pathology. *Prog. Brain Res.*, **162**, 245-273.
- Smith, J., Ladi, E., Mayer-Proschel, M. and Noble, M. (2000): Redox state is a central modulator of the balance between self-renewal and differentiation in a dividing glial precursor cell. *Proceedings of the National Academy of Sciences of the United States of America*, **97**, 10032-10037.
- Sripetchwandee, J., Sanit, J., Chattipakorn, N. and Chattipakorn, S.C. (2013): Mitochondrial calcium uniporter blocker effectively prevents brain mitochondrial dysfunction caused by iron overload. *Life Sci.*, **92**, 298-304.
- Topchiy, E., Panzhinskiy, E., Griffin, W.S., Barger, S.W., Das, M. and Zawada, W.M. (2013): Nox4-generated superoxide drives angiotensin II-induced neural stem cell proliferation. *Dev. Neurosci.*, **35**, 293-305.
- Touati, D. (2000): Iron and oxidative stress in bacteria. *Arch. Biochem. Biophys.*, **373**, 1-6.
- Tyagi, P.K., Janowska, I., Cretu, O., Pham-Huu, C. and Banhart, F. (2011): Catalytic action of gold and copper crystals in the growth of carbon nanotubes. *J. Nanosci. Nanotechnol.*, **11**, 3609-3615.
- Xu, L.J., Zhao, J.X., Zhang, T., Ren, G.G. and Yang, Z. (2009): *In vitro* study on influence of nano particles of CuO on CA1 pyramidal neurons of rat hippocampus potassium currents. *Environ. Toxicol.*, **24**, 211-217.
- Yazyev, O.V. and Pasquarello, A. (2008): Effect of metal elements in catalytic growth of carbon nanotubes. *Phys. Rev. Lett.*, **100**, 156102.
- Yim, M.B., Chae, H.Z., Rhee, S.G., Chock, P.B. and Stadtman, E.R. (1994): On the protective mechanism of the thiol-specific antioxidant enzyme against the oxidative damage of biomacromolecules. *J. Biol. Chem.*, **269**, 1621-1626.
- Zager, R.A. and Burkhart, K.M. (1998): Differential effects of glutathione and cysteine on Fe²⁺, Fe³⁺, H₂O₂ and myoglobin-induced proximal tubular cell attack. *Kidney Int.*, **53**, 1661-1672.
- Zhang, Z., Teruya, K., Eto, H. and Shirahata, S. (2011): Fucoidan extract induces apoptosis in MCF-7 cells via a mechanism involving the ROS-dependent JNK activation and mitochondria-mediated pathways. *PLoS One*, **6**, e27441.