









10% O<sub>2</sub> Z-VAD-FMK  
+ BHT

10% O<sub>2</sub> Y-27632

10% O<sub>2</sub> HGF

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + Y-27632



10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + HGF

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + Y-27632 +  
HGF

10% O<sub>2</sub> Control

20% O<sub>2</sub> Blocked  
Control









Supp. Fig. 1: Morphologies observed in the different culture conditions 24h after seeding.

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT

10% O<sub>2</sub> Y-27632

10% O<sub>2</sub> HGF

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + Y-27632



10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + HGF

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + Y-27632 +  
HGF

10% O<sub>2</sub> Control

20% O<sub>2</sub> Blocked  
Control









Supp. Fig. 2: Morphologies observed in the different culture conditions 48h after seeding.

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT

10% O<sub>2</sub> Y-27632

10% O<sub>2</sub> HGF

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + Y-27632



10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + HGF

10% O<sub>2</sub> Z-VAD-FMK  
+ BHT + Y-27632 +  
HGF

10% O<sub>2</sub> Control

20% O<sub>2</sub> Blocked  
Control

Supp. Fig. 3: Morphologies observed in the different culture conditions 72h after seeding.